

# Geometry of Fourfolds with an Admissible K3 Subcategory

Laura Pertusi

Dipartimento di Matematica F. Enriques, Università degli Studi di Milano

Advisor: Paolo Stellari

## Setting

A **K3 category** is a triangulated category whose Serre functor is the shift  $(-)[2]$  and with the same Hochschild cohomology of a K3 surface. Cubic fourfolds and Gushel-Mukai fourfolds have a semiorthogonal decomposition of their derived category of coherent sheaves given by exceptional objects and an admissible K3 category. This allows us to study:

- Fourier-Mukai partners of cubic fourfolds;
- the double EPW sextic associated to a GM fourfold as a moduli space of twisted sheaves on a K3 surface.

**Motivation:** K3 categories simplify the study of moduli problems over cubic or GM fourfolds.

## Fourier-Mukai partners of cubic fourfolds

A cubic fourfold  $X$  is a smooth cubic hypersurface in  $\mathbb{P}_{\mathbb{C}}^5$ .

### Semiorthogonal decomposition

$$D^b(X) = \langle \mathcal{A}_X, \mathcal{O}_X, \mathcal{O}_X(1), \mathcal{O}_X(2) \rangle$$

where  $\mathcal{A}_X$  is a K3 category (Kuznetsov).

**Definition:** A cubic fourfold  $X'$  is a **FM partner** of  $X$  if there is an equivalence  $\mathcal{A}_X \xrightarrow{\sim} \mathcal{A}_{X'}$  of Fourier-Mukai type, i.e. there exists  $K \in D^b(X \times X')$  such that

$$\Phi : D^b(X) \rightarrow \mathcal{A}_X \xrightarrow{\sim} \mathcal{A}_{X'} \rightarrow D^b(X')$$

$$\Phi(-) \cong R p_{X'*}(K \otimes^L L p_X^*(-)).$$

Consistently with the analogy to K3 surfaces we have:

**Theorem:** ([1]) The number of isomorphism classes of FM partners  $\#FM(X)$  of  $X$  is finite.

## Question

Are there examples of cubic fourfolds with a prescribed number of non isomorphic FM partners?

**Answer:** Consider **general** cubic fourfolds of discriminant  $d$  with a **Hodge-associated (twisted) K3 surface**  $(S, \alpha) \Leftrightarrow$  numerical condition on the discriminant.

## Untwisted case (Hassett)

$$4 \nmid d, 9 \nmid d, p \nmid d \forall \text{ prime } p \equiv 2 \pmod{3} \quad (*)$$

## Twisted case (Huybrechts)

$$n_i \equiv 0 \pmod{2} \forall \text{ prime } p_i \equiv 2 \pmod{3} \text{ in } 2d = \prod p_i^{n_i} \quad (*')$$

## Theorem 1

Let  $d > 6$ ,  $d \equiv 0, 2 \pmod{6}$  satisfying  $(*)'$  and let  $h$  be the number of distinct odd primes in prime factorization of  $d/\text{ord}(\alpha)$ . Let  $X$  be a general element in  $\mathcal{C}_d$ .

**Untwisted case:** If  $d$  satisfies  $(*)$ , then

$$\#FM(X) = \begin{cases} 2^{h-1}, & \text{if } d \equiv 2 \pmod{6} \text{ and } h > 1; \\ 2^{h-2}, & \text{if } d \equiv 0 \pmod{6} \text{ and } h > 2; \\ 1, & \text{otherwise.} \end{cases}$$

**Twisted case:** We get a lower bound to  $\#FM(X)$ , depending on  $h$  and  $\varphi(\alpha)$ .

## Tool: Mukai lattice for $\mathcal{A}_X$

$$\tilde{H}(\mathcal{A}_X, \mathbb{Z}) = \{ \kappa \in K_{\text{top}}(X) : \chi([\mathcal{O}_X(i)], \kappa) = 0, \forall i = 0, 1, 2 \}.$$

**Theorem:** ([1]) For general  $X \in \mathcal{C}_d$ ,  $\mathcal{A}_X \xrightarrow{\sim} \mathcal{A}_{X'}$  of FM type iff their Mukai lattices are Hodge isometric.

## Gushel-Mukai fourfolds

A GM fourfold is a smooth dimensionally transverse intersection

$$X = \text{CG}(2, V_5) \cap \mathbb{P}(W) \cap Q,$$

where  $Q$  is a quadric hypersurface in  $\mathbb{P}(W) \cong \mathbb{P}^8 \subset \mathbb{P}(\wedge^2 V_5 \oplus \mathbb{C})$ .

### Semiorthogonal decomposition

$$D^b(X) = \langle \mathcal{A}_X, \mathcal{O}_X, \mathcal{U}_X^*, \mathcal{O}_X(1), \mathcal{U}_X^*(1) \rangle$$

where  $\mathcal{A}_X$  is a K3 category (Kuznetsov, Perry).

**Lagrangian data:**  $X$  defines a triple  $(V_6, V_5, A)$ , where  $A \subset \wedge^3 V_6$  is Lagrangian without decomposable vectors. Viceversa, it is possible to recover the GM fourfold from such a data (Debarre, Kuznetsov).

$\rightsquigarrow$  EPW stratification in  $Y_A^{\geq 3} \subset Y_A^{\geq 2} \subset Y_A^{\geq 1} \subset \mathbb{P}(V_6)$  and **EPW sextic** hypersurface  $Y_A := Y_A^{\geq 1}$ .

## Associated double EPW sextic

We consider the double cover of the EPW sextic  $Y_A$  branched over  $Y_A^{\geq 2}$  associated to a GM fourfold  $X$ . Assume that  $\tilde{Y}_A$  is smooth  $\Leftrightarrow Y_A^{\geq 3} = \emptyset$ .

## Aim

To study  $\tilde{Y}_A$  as a moduli space of (twisted) stable sheaves on a K3 surface.

## Facts (Debarre, Iliev, Manivel):

- Period points of **special** GM fourfolds form divisors in the period domain identified by the discriminant  $d$ .
- Hodge-associated K3 surface  $\Leftrightarrow 8 \nmid d$  and the only odd primes which divide  $d$  are  $\equiv 1 \pmod{4}$   $(\dagger)$

Assume that  $X$  has discriminant  $d$ .

## Theorem 2 (untwisted case)

If  $d$  satisfies  $(\dagger)$ , then  $\tilde{Y}_A$  is birational to a moduli space of stable sheaves on a K3 surface  $S$ .

The converse holds for general  $X$  and for non general  $X$  whose period point is in a divisor with discriminant  $d \equiv 2$  or  $4 \pmod{8}$ .

**Remark:** There are examples of GM fourfolds with  $\text{rank}(H^{2,2}(X, \mathbb{Z})) = 4$  and period point only in divisors with discriminants  $\equiv 0 \pmod{8}$ , having  $\tilde{Y}_A$  birational to a moduli space of sheaves on a K3 surface, **but** which cannot have a Hodge-associated K3 surface.

## Steps:

- Mukai lattice  $\tilde{H}(\mathcal{A}_X, \mathbb{Z})$

$$\langle \lambda_1, \lambda_2 \rangle^\perp \cong H^4(X, \mathbb{Z})_{\text{van}}$$

- Relate condition  $(\dagger)$  with existence of a primitively embedded  $U = (\mathbb{Z}^2, \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix})$  in the algebraic part of  $\tilde{H}(\mathcal{A}_X, \mathbb{Z})$ .
- There is a primitive embedding of  $H^2(\tilde{Y}_A, \mathbb{Z})$  in  $\tilde{H}(\mathcal{A}_X, \mathbb{Z}) \rightsquigarrow$  we apply Addington's result.

## Theorem 2 (twisted case)

There is a Hodge isometry  $\tilde{H}(\mathcal{A}_X, \mathbb{Z}) \cong \tilde{H}(S, \alpha, \mathbb{Z})$  where  $(S, \alpha)$  is a twisted K3 surface iff

$$d = \prod_i p_i^{n_i} \text{ with } n_i \equiv 0 \pmod{2} \text{ for } p_i \equiv 3 \pmod{4} \quad (\dagger')$$

$\tilde{Y}_A$  is birational to a moduli space of twisted stable sheaves on a K3 surface  $S$  if and only if  $d$  satisfies  $(\dagger')$ .

## Theorem 3

$\tilde{Y}_A$  is birational to the Hilbert scheme  $S^{[2]}$  on a K3 surface  $S$  iff  $d$  satisfies

$$a^2 d = 2n^2 + 2 \quad \text{for } a, n \in \mathbb{Z}.$$

## Stability conditions on $\mathcal{A}_X$ (work in progress)

**Property:** ([2]) If  $X$  is a generic GM fourfold, then the restriction to a hyperplane  $\mathbb{P}(V_4) \subset \mathbb{P}(V_5)$  of the first conic fibration  $\rho$  is flat and smooth.

$$\begin{array}{ccc} \tilde{X} & \longrightarrow & \mathbb{P}_X(\mathcal{U}_X) \\ \sigma \swarrow & \tilde{\rho} \downarrow & \downarrow \rho \\ X & \longleftarrow & \mathbb{P}(V_4) \longrightarrow \mathbb{P}(V_5) \end{array}$$

$\tilde{X} = \text{Bl}_E(X)$ ,  $E = G(2, V_4) \cap Q$ .

**Idea:** Use  $\tilde{\rho}$  to induce stability conditions on  $\mathcal{A}_X$  from  $D^b(\mathbb{P}^3, \mathcal{B}_0)$ ,  $\mathcal{B}_0 =$  even part of associated Clifford algebra.

## References

- [1] D. Huybrechts, *The K3 category of a cubic fourfold*, *Compositio Mathematica* **153** (2017), 586-620.
- [2] M. Ornaghi, L. Pertusi, *Voevodsky's conjecture for cubic fourfolds and Gushel-Mukai fourfolds via noncommutative K3 surfaces*, arXiv:1703.10844.
- [3] L. Pertusi, *Fourier-Mukai partners for general special cubic fourfolds*, arXiv:1611.06687
- [4] L. Pertusi, *On the double EPW sextic associated to a Gushel-Mukai fourfold*, in preparation.